

# UNCLASSIFIED

AD NUMBER
AD866109
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; JAN 1970. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Release Branch/TID, Frederick, MD 21701.
AUTHORITY
BDRL D/A ltr, 29 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

866109

## TECHNICAL MANUSCRIPT 580

# EFFECT OF RECIPIENT CELL CONCENTRATION ON TRANSFECTION WITH BACTERIOPHAGE DNA

Darrel D. Gwinn  
William D. Lawton

### STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch/TID, Frederick, Maryland 21701

JANUARY 1970

DEPARTMENT OF THE ARMY

Fort Detrick  
Frederick, Maryland

Reproduced by the  
CLEARINGHOUSE  
for Federal Scientific & Technical  
Information Springfield Va. 22151

DDC  
RECEIVED  
MAR 10 1970  
REGULATED  
C

49584

16

ACCESSION FOR		
CESTI	WHITE SECTION <input type="checkbox"/>	
DUG	RWY SECTION <input checked="" type="checkbox"/>	
ANNOUNCED <input type="checkbox"/>		
DISSEMINATION		
DISTRIBUTION/AVAILABILITY CODES		
QST.	AVAIL.	FOR SPECIAL
2		

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

#### DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

#### DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DEPARTMENT OF THE ARMY  
Fort Detrick  
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 580

EFFECT OF RECIPIENT CELL CONCENTRATION ON TRANSFECTION  
WITH BACTERIOPHAGE DNA

Darrel D. Gwinn

William D. Lawton

Medical Bacteriology Division  
BIOLOGICAL SCIENCES LABORATORIES

Project 1B061102B71A

January 1970

#### ACKNOWLEDGMENT

We acknowledge the able technical assistance of Raymond G. Gregoire.

#### ABSTRACT

Transfection of Bacillus subtilis 168 with low concentrations of SP-18 phage DNA (molecular weight =  $1 \times 10^8$ ) was enhanced by using recipient cells at a concentration of  $2 \times 10^7$ /ml rather than the usual  $2 \times 10^8$ /ml. At low DNA concentrations, the slope of the dose-response curve was  $>1$  at the high recipient cell concentration, but the slope was  $= 1$  at the low recipient cell concentration. A similar effect was shown with SP-82 DNA,  $\phi 1$  DNA, and  $\phi 25$  DNA, all having molecular weights of approximately  $1 \times 10^8$ . However,  $\phi 29$  DNA (molecular weight =  $2.5 \times 10^7$ ) gave a first-order dose-response curve at both high and low recipient cell concentrations. We interpreted our observations as Kelly hypothesized for the preservation of marker linkage in transformation; i.e., the lower concentration of recipient cells resulted in fewer DNA molecules being fragmented as a result of attaching to two cells. This interpretation of transfection enhancement is different from the previous explanation as an inhibition of an intracellular inactivation process.

# CONTENTS

Acknowledgment . . . . .	2
Abstract . . . . .	2
I. INTRODUCTION . . . . .	5
II. MATERIALS AND METHODS . . . . .	5
III. RESULTS . . . . .	6
IV. DISCUSSION . . . . .	13
Literature Cited . . . . .	15
Distribution List . . . . .	17
DD Form 1473 . . . . .	19

# FIGURES

1. Effect of SP-18 DNA Concentration on Formation of Infectious Centers . . . . .	7
2. Effect of Recipient Cell Concentration on Formation of Infectious Centers . . . . .	7
3. Effect of SP-18 DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers . . . . .	8
4. Effect of SP-18 DNA Concentration and Washed Recipient Cell Concentration on Formation of Infectious Centers . . . . .	9
5. Effect of SP-82 DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers . . . . .	9
6. Effect of Ø29 DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers . . . . .	11
7. Effect of Ø25 DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers . . . . .	12
8. Effect of Ø1 DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers . . . . .	12

## I. INTRODUCTION\*

In our studies to determine why the infectivity of SP-18 phage DNA was nonlinear at low DNA concentrations, we observed that dilution of the recipient cells greatly increased the number of transfectants obtained with low concentrations of SP-18 DNA.

Our observation was similar to that reported by Kelly<sup>1</sup> in which the use of dilute suspensions of competent cells preserved the linkage of various markers in transformation. We felt it worthwhile to expand this observation and to study the use of diluted recipient cells for transfection by DNA of various bacteriophages.

## II. MATERIALS AND METHODS

Phage SP-18 was isolated from soil by Dr. Ivan D. Goldberg in this laboratory. Phages  $\phi$ 1,  $\phi$ 25, and  $\phi$ 29 were described by Kelly and Spizizen,<sup>2</sup> and phage SP-82 was described by Green.<sup>3</sup> All phage suspensions were assayed on phage assay agar (PA) seeded with  $10^8$  *Bacillus subtilis* 168 (*ind*<sup>-</sup>) spores as described by Thorne.<sup>4</sup> Cultures of *B. subtilis* 168 (*ind*<sup>-</sup>) were used to prepare SP-18,  $\phi$ 1,  $\phi$ 25, and  $\phi$ 29 in TY broth<sup>5</sup> and SP-82 in H broth.<sup>6</sup>

Phage DNA was extracted by a modification of the phenol technique of Mandell and Hershey.<sup>7</sup> The DNA was dialyzed for 22 hours at 5 C against 0.15 M NaCl and 0.015 M citrate (SSC) at pH 8.0 to remove the phenol and was stored at 5 C. DNA was determined by the method of Burton.<sup>8</sup> All DNA preparations were tested for contamination with bacteria or viable phage by spreading samples on nutrient agar and by plating samples (after treatment with 50  $\mu$ g of deoxyribonuclease) in PA agar seeded with  $10^8$  *B. subtilis* 168 (*ind*<sup>-</sup>) spores. Serial twofold dilutions of the DNA preparations were made in SSC.

Competent cells of *B. subtilis* 168 (*ind*<sup>-</sup>) were grown essentially by the procedure of Anagnostopoulos and Spizizen.<sup>9</sup> Transfections were carried out in 18- by 100-mm tubes containing 1 ml of recipient culture (approximately  $2 \times 10^8$  or  $2 \times 10^7$  cells) and 0.1 ml of DNA. The tubes were incubated for 45 minutes in a slanted position on a reciprocal shaker at 37 C. Pancreatic deoxyribonuclease (IX crystallized, Worthington) was added (50  $\mu$ g in 0.05 ml) and incubation was continued for 5 minutes. Infectious centers were scored against 168 (*ind*<sup>-</sup>) by the soft agar overlay method in PA agar.

---

\* This report should not be used as a literature citation in material to be published in the open literature. Readers interested in referencing the information contained herein should contact the senior author to ascertain when and where it may appear in citable form.

### III. RESULTS

SP-18 DNA showed a first-order relationship between DNA concentration and infectivity in the region from approximately 2 to 40  $\mu\text{g/ml}$ , but at DNA concentrations below 2  $\mu\text{g/ml}$  the infectivity was greater than first order (Fig. 1). To determine the relationship between infectivity of SP-18 DNA and cell concentration, we made serial twofold dilutions of the recipient cells in the presence of DNA at 10  $\mu\text{g/ml}$  and 1.25  $\mu\text{g/ml}$  (Fig. 2). At 1.25  $\mu\text{g DNA/ml}$ , maximum transfection was demonstrated with  $2 \times 10^7$  cells/ml; more or fewer cells resulted in fewer transfectants/ml.

The effect of SP-18 DNA concentration on infectious center formation at recipient cell concentrations of approximately  $2 \times 10^8$  and  $2 \times 10^7$  cells/ml is shown in Figure 3. The dose-response curve at the high cell concentration was the same as that in Figure 1, but at the low cell concentration, the dose-response curve remained first-order in the region from about 0.2 to 2  $\mu\text{g/ml}$ . This resulted in a higher percentage transfection with a cell concentration of approximately  $2 \times 10^7$  cells/ml; e.g., at a DNA concentration of 0.2  $\mu\text{g/ml}$ , the transfection obtained with  $2 \times 10^7$  cells/ml was more than 300% that obtained with  $2 \times 10^8$  cells/ml.

The increase in percentage transfection obtained with diluted competent 168 (*ind*<sup>-</sup>) cells could be explained if, in diluting the cells, we also diluted a competence inhibitor or extracellular nuclease. To test this possibility, competent 168 (*ind*<sup>-</sup>) cells were washed to get rid of any extracellular competence inhibitor or nuclease. However, the dose-response curves from washed cells (Fig. 4) were the same as those obtained from nonwashed cells (Fig. 3). An alternative explanation for our observed increase in percentage transfection with diluted recipient cells was the hypothesis presented by Kelly<sup>1</sup> that a DNA molecule might adsorb to more than one bacterial cell at the same time, resulting in fragmentation of the molecule. Dilution of the competent cells would lower the chance of a DNA molecule adsorbing to more than one cell, lowering fragmentation, and thereby increasing transfection.

To determine if the use of diluted recipient cells affected transfection with other phage DNA, we tested phage SP-82, whose DNA has a molecular weight ( $1.3 \times 10^6$ ) similar to that of SP-18 DNA ( $1.0 \times 10^6$ ). SP-82 DNA does not give a first-order dose-response curve in any region of DNA concentrations unless an enhancement mechanism is used.<sup>10</sup> However, when a recipient cell concentration of approximately  $2 \times 10^7$  was used, we obtained a first-order dose response with SP-82 DNA (Fig. 5). If large DNA molecules were fragmented by adsorption to more than one recipient cell, we would expect smaller DNA molecules to be less affected. To test this, we measured transfection with  $\phi 29$  DNA, which has a molecular weight of  $2.5 \times 10^7$ . The  $\phi 29$  DNA gave a first-order relationship between DNA concentration and infectious-center formation at the same DNA concentrations for both high and






FIGURE 1. Effect of SP-18 DNA Concentration on Formation of Infectious Centers.  $\circ$  = approximately  $2 \times 10^8$  recipient cells/ml.




FIGURE 2. Effect of Recipient Cell Concentration on Formation of Infectious Centers.  $\circ$  = 10  $\mu$ g SP-18 DNA/ml,  $\Delta$  = 1.25  $\mu$ g SP-18 DNA/ml.




FIGURE 3. Effect of SP-18 DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers. ○ = approximately  $2 \times 10^7$  recipient cells/ml, △ = approximately  $2 \times 10^8$  recipient cells/ml.




FIGURE 4. Effect of SP-18 DNA Concentration and Washed Recipient Cell Concentration on Formation of Infectious Centers. The competent 168 (ind<sup>-</sup>) cells were washed two times in competence media and resuspended in competence media to their original titer. O = approximately  $2 \times 10^7$  recipient cells/ml,  $\Delta$  = approximately  $2 \times 10^8$  recipient cells/ml.




FIGURE 5. Effect of SP-82 DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers. O = approximately  $2 \times 10^7$  recipient cells/ml,  $\Delta$  = approximately  $2 \times 10^8$  recipient cells/ml.

low recipient cell concentration (Fig. 6). Our interpretation was that transfection with low molecular weight  $\phi$ 29 DNA was not affected by fragmentation of the DNA molecules resulting from adsorption to more than one recipient cell. Concentrations of  $\phi$ 29 DNA lower than those shown in Figure 6 gave a dose-response curve with a slope  $>1$  at both high and low recipient cell concentrations.

To show that the first-order dose response obtained with low concentrations of recipient cells was not unique to SP-18 DNA and SP-82 DNA, we transfected high and low recipient cell concentrations with  $\phi$ 1 DNA and  $\phi$ 25 DNA, both having molecular weights of  $1 \times 10^8$ . The results (Fig. 7 and 8) supported our previous observations with high molecular weight DNA; we obtained a first-order dose-response curve with the low recipient cell concentrations but not with the high recipient cell concentration.

When phage DNA concentrations higher than those shown were added to high recipient cell concentrations, we observed a flattening of the dose-response curve or saturation of the system. It must be emphasized that a first-order dose response from SP-82,  $\phi$ 1, and  $\phi$ 25 DNA was obtained only with low recipient cell concentrations.




FIGURE 6. Effect of  $\phi 29$  DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers.  $\circ$  = approximately  $2 \times 10^7$  recipient cells/ml,  $\Delta$  = approximately  $2 \times 10^8$  recipient cells/ml.




FIGURE 8. Effect of  $\phi 1$  DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers. O = approximately  $2 \times 10^7$  recipient cells/ml,  $\Delta$  = approximately  $2 \times 10^8$  recipient cells/ml.




FIGURE 7. Effect of  $\phi 25$  DNA Concentration and Recipient Cell Concentration on Formation of Infectious Centers. O = approximately  $2 \times 10^7$  recipient cells/ml,  $\Delta$  = approximately  $2 \times 10^8$  recipient cells/ml.

#### IV. DISCUSSION

Our observations can be interpreted to support Kelly's hypothesis<sup>1</sup> that "at low concentrations of high molecular weight DNA, molecules normally attach to and are integrated into more than one recipient cell."

The dose-response curves from SP-82, Ø1, and Ø25 DNA using high recipient cell concentrations went from saturation to curves with slopes  $>1$ . There was no intermediate DNA concentration where a first-order dose response was obtained.

The first-order dose-response curves from SP-18 DNA, SP-82 DNA, Ø1 DNA, and Ø25 DNA that we observed at low concentrations of recipient cells change to greater than first order at the lowest DNA concentrations. The most probable explanation for this change is loss of contact with the recipient cells by the dilute, high molecular weight DNA, resulting in fewer infections being initiated.

SP-18 DNA gave a first-order dose response at both high and low recipient cell concentrations, but not at the same DNA concentration. SP-82 DNA, Ø1 DNA, and Ø25 DNA gave first-order dose responses only at low recipient cell concentrations. The genomes of these three phages may have weak points making them more susceptible to fragmentation.

SP-82 DNA, which normally shows a dose-response curve with a slope  $>1$ , has been reported to show a dose-response curve with a slope of one under certain conditions. Green<sup>11</sup> obtained a first-order curve with SP-82 by the use of helper phage, and he hypothesized that preinfection with mutant helper phage inhibited the intracellular inactivation process that he felt was responsible for the dose-response curve with a slope  $>1$ .

Epstein<sup>12</sup> enhanced SP-82 transfection by UV-irradiation of the recipient cells, and Epstein and Mahler<sup>10</sup> enhanced SP-82 transfection by pretransfection exposure of competent cells to UV-irradiated DNA.

LITERATURE CITED

1. Kelly, M.S. 1967. The causes of instability of linkage in transformation of Bacillus subtilis. Mol. Gen. Genet. 99:350-361.
2. Reilly, B.E.; Spizizen, J. 1965. Bacteriophage deoxyribonucleate infection of competent Bacillus subtilis. J. Bacteriol. 89:782-790.
3. Green, D.M. 1964. Infectivity of DNA isolated from Bacillus subtilis bacteriophage, SP-82. J. Mol. Biol. 10:438-451.
4. Thorne, C.B. 1962. Transduction in Bacillus subtilis. J. Bacteriol. 83:106-111.
5. Romig, W.R. 1962. Infection of Bacillus subtilis with phenol-extracted bacteriophages. Virology 16:452-459.
6. Chase, M.; Doermann, A.H. 1958. High negative interference over short segments of the genetic structure of bacteriophage T<sub>4</sub>. Genetics 43:332-353.
7. Mandell, J.D.; Hershey, A.D. 1960. A fractionating column for analysis of nucleic acids. Anal. Biochem. 1:66-77.
8. Burton, K. 1956. A study of the conditions and mechanism of the diphenylamine reaction for the colorimetric estimation of desoxyribonucleic acid. Biochem. J. 62:315-322.
9. Anagnostopoulos, C.; Spizizen, J. 1961. Requirements for transformation in Bacillus subtilis. J. Bacteriol. 81:741-746.
10. Epstein, H.T.; Mahler, I. 1968. Mechanisms of enhancement of SP-82 transfection. J. Virol. 2:710-715.
11. Green, D.M. 1966. Intracellular inactivation of infective SP-82 bacteriophage DNA. J. Mol. Biol. 22:1-14.
12. Epstein, H.T. 1967. Transfection enhancement by ultraviolet light. Biochem. Biophys. Res. Commun. 27:258-262.



Unclassified  
Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Department of the Army Fort Detrick, Frederick, Maryland, 21701		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE EFFECT OF RECIPIENT CELL CONCENTRATION ON TRANSFECTION WITH BACTERIOPHAGE DNA		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name) Darrel D. Gwinn William D. Lawton		
6. REPORT DATE January 1970	7a. TOTAL NO. OF PAGES 19	7b. NO. OF REFS 12
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 580
b. PROJECT NO. 1B061102B71A		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) CMS 6642 SMUFD-AE-T 49584
c. Task-Work Unit 01-022		
d. DD 1498 Agency Accession DA OL 0193		
10. DISTRIBUTION STATEMENT Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES Medical Bacteriology Division		12. SPONSORING MILITARY ACTIVITY Department of the Army Fort Detrick, Frederick, Maryland, 21701
13. ABSTRACT Transfection of <u>Bacillus subtilis</u> 168 with low concentrations of SP-18 phage DNA (molecular weight = $1 \times 10^8$ ) was enhanced by using recipient cells at a concentration of $2 \times 10^7$ /ml rather than the usual $2 \times 10^8$ /ml. At low DNA concentrations, the slope of the dose-response curve was $>1$ at the high recipient cell concentration, but the slope was $= 1$ at the low recipient cell concentration. A similar effect was shown with SP-82 DNA, $\phi 1$ DNA, and $\phi 25$ DNA, all having molecular weights of approximately $1 \times 10^8$ . However, $\phi 29$ DNA (molecular weight = $2.5 \times 10^7$ ) gave a first-order dose-response curve at both high and low recipient cell concentrations. We interpreted our observations as Kelly hypothesized for the preservation of marker linkage in transformation; i.e., the lower concentration of recipient cells resulted in fewer DNA molecules being fragmented as a result of attaching to two cells. This interpretation of transfection enhancement is different from the previous explanation as an inhibition of an intracellular inactivation process.		
14. Key Words Transfection DNA Cell concentration Dose-response curve Bacteriophage <u>Bacillus subtilis</u> Infectivity		